



THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

# High-sensitivity cardiac troponin concentrations at presentation in patients with ST-segment elevation myocardial infarction

### Citation for published version:

Wereski, R, Chapman, A, Lee, K, Smith, SW, Lowe, DJ, Gray, A & Mills, N 2020, 'High-sensitivity cardiac troponin concentrations at presentation in patients with ST-segment elevation myocardial infarction', *JAMA Cardiology*, vol. 5, no. 11, pp. 1302–1304. <https://doi.org/10.1001/jamacardio.2020.2867>

### Digital Object Identifier (DOI):

[10.1001/jamacardio.2020.2867](https://doi.org/10.1001/jamacardio.2020.2867)

### Link:

[Link to publication record in Edinburgh Research Explorer](#)

### Document Version:

Publisher's PDF, also known as Version of record

### Published In:

JAMA Cardiology

### General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

### Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



# Letters

## RESEARCH LETTER


### High-Sensitivity Cardiac Troponin Concentrations at Presentation in Patients With ST-Segment Elevation Myocardial Infarction

The introduction of high-sensitivity cardiac troponin testing into clinical practice has transformed the assessment of patients with suspected acute coronary syndrome in the emergency department.<sup>1</sup> Most patients can be discharged using accelerated diagnostic pathways that do not require hospital admission for peak cardiac troponin testing.<sup>2</sup> These pathways are not recommended for patients with ST-segment elevation on the electrocardiogram,<sup>3,4</sup> but given that interpretation

is dependent on experience, there is a risk patients could be inappropriately assessed.

**Methods** | Between June 2013 and March 2016, consecutive patients with suspected acute coronary syndrome were recruited across 10 hospitals in the High-Sensitivity Troponin in the Evaluation of Patients With Acute Coronary Syndrome (High-STEACS) cluster randomized clinical trial.<sup>5</sup> High-sensitivity cardiac troponin I was measured using the Abbott ARCHITECT STAT assay (Abbott Laboratories), which has a limit of detection of 1.2 ng/L, and a 99th percentile upper reference limit of 34 ng/L in men and 16 ng/L in women. The index diagnosis was independently adjudicated by 2 physicians on

**Figure. High-Sensitivity Cardiac Troponin I (hs-TnI) Concentrations in Consecutive Patients With ST-Segment Elevation Myocardial Infarction Stratified by Troponin Concentration at Presentation**



Patients were stratified by time of onset of symptoms (<2 hours and ≥2 hours) and according to cardiac troponin concentration at presentation. Individual patient concentrations (less than the rule-out threshold of <5 ng/L [green; to convert to micrograms per liter, multiply by 1], 5 ng/L to the 99th percentile

diagnostic threshold [yellow], 99th percentile to 52 ng/L rule-in threshold for the ESC 0/1 hours pathway [orange], and >52 ng/L [red]) are shown with box and whisker distribution and probability density plots. Peak concentration is the highest troponin concentration obtained on serial sampling.

review of all clinical information, according to the Universal Definition of Myocardial Infarction.<sup>6</sup> Patients with type 1 ST-segment elevation myocardial infarction (STEMI) were stratified according to cardiac troponin concentration at presentation using a validated risk-stratification threshold (5 ng/L),<sup>1</sup> the sex-specific 99th percentile, and the European Society of Cardiology (ESC) 0 of 1 hour pathway rule-in threshold (52 ng/L).<sup>4</sup> Posterior STEMI was defined as those with STEMI and an acute occlusion of the circumflex, obtuse marginal, or posterior left ventricular artery on angiography. Time from symptom onset was recorded prospectively by attending clinicians. The trial was approved by the National Health Service Scotland A Research Ethics Committee, the Public Benefit and Privacy Panel for Health and Social Care, and by each local National Health Service Health Board. Because randomization was at the hospital level, individual patient consent was not sought.<sup>5</sup> Comparisons between groups were performed using the  $\chi^2$  test for categorical variables and an unpaired *t* test or the Kruskal-Wallis test for continuous variables. Statistical analysis was performed using R, version 3.6.1 (R Foundation).

**Results** | The trial enrolled 48 282 consecutive patients, of whom 925 had an adjudicated diagnosis of STEMI (67.8% men [*n* = 627 of 925]; mean [SD] age, 65 [14] years). At presentation, the median troponin concentration was 196 ng/L (interquartile range [IQR], 46-21 611 ng/L), with 2.2% (*n* = 20 of 925) and 14.4% of patients (*n* = 133 of 925) having concentrations less than 5 ng/L and the 99th percentile, respectively (Figure). Just 73.2% of patients (*n* = 677 of 925) had troponin concentrations greater than the rule-in threshold of 52 ng/L. Patients presenting within 2 hours of symptom onset (23.4%; 216 of 809) had lower troponin concentrations (96 ng/L; IQR, 26-494 ng/L vs 294 ng/L; IQR, 59-3042 ng/L; *P* < .001) and were more likely to have concentrations at less than the 99th percentile (26.4% [*n* = 57 of 216] vs 14.1% [*n* = 95 of 674]; *P* < .001), compared with those presenting later. Posterior STEMI was more common in patients presenting with troponin at less than the 99th percentile (18.1% [*n* = 26 of 144] vs 9.8% [*n* = 61 of 618]; *P* = .008).

**Discussion** | Despite significant advances in the sensitivity of cardiac troponin testing, more than 1 in 4 patients with STEMI have troponin concentrations at less than the ESC-recommended rule-in threshold at presentation. Patients presenting within 2 hours were more likely to have a troponin concentration at less than the 99th percentile; however, even in those who presented later, 1 in 6 had troponin concentrations at less than the diagnostic threshold. During myocardial infarction, abrupt coronary occlusion may prevent the release of troponin into the circulation until reperfusion has occurred. Our observations are an important reminder of the limited role of troponin testing in the early assessment of patients with ST-segment elevation. Where clinical suspicion is high, troponin concentrations within the reference range should not delay the initiation of therapeutic agents or urgent coronary angiography. This is particularly relevant in patients with electrocardiographic changes suspicious of posterior myocardial infarction, but our findings are relevant to a wider group of patients with conduction abnormalities, such as bundle branch block

or ventricular pacing, where interpretation of the electrocardiogram is challenging.

Ryan Wereski, MD  
 Andrew R. Chapman, MD, PhD  
 Ken K. Lee, MD  
 Stephen W. Smith, MD  
 David J. Lowe, MD  
 Alasdair Gray, MD  
 Nicholas L. Mills, MD, PhD

**Author Affiliations:** BHF Centre for Cardiovascular Science, University of Edinburgh, Edinburgh, Scotland (Wereski, Chapman, Lee, Mills); University of Minnesota, Emergency Medicine, Minneapolis (Smith); University of Glasgow, School of Medicine, Glasgow, Scotland (Lowe); Royal Infirmary of Edinburgh, Emergency Medicine Research Group, Edinburgh, Scotland (Gray).

**Corresponding Author:** Nicholas L. Mills, MD, PhD, BHF/University Centre for Cardiovascular Science, the University of Edinburgh, Edinburgh EH16 4SA, Scotland (nick.mills@ed.ac.uk).

**Accepted for Publication:** May 29, 2020.

**Published Online:** August 12, 2020. doi:10.1001/jamacardio.2020.2867

**Correction:** This article was corrected on September 23, 2020, to fix errors in the Figure.

**Open Access:** This is an open access article distributed under the terms of the CC-BY License. © 2020 Wereski R et al. *JAMA Cardiology*.

**Author Contributions:** Drs Wereski and Chapman had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

**Concept and design:** Wereski, Chapman, Lee, Smith, Lowe, Mills.

**Acquisition, analysis, or interpretation of data:** Wereski, Chapman, Gray, Mills.

**Drafting of the manuscript:** Wereski, Chapman, Gray, Mills.

**Critical revision of the manuscript for important intellectual content:** Chapman, Lee, Smith, Lowe, Mills.

**Statistical analysis:** Wereski, Chapman, Lee.

**Obtained funding:** Mills.

**Administrative, technical, or material support:** Chapman, Mills.

**Supervision:** Chapman, Smith, Lowe, Mills.

**Conflict of Interest Disclosures:** Dr Mills has received honoraria or consultancy from Abbott Diagnostics, Roche Diagnostics, Siemens Healthineers, and LumiraDx. No other disclosures were reported.

**Funding/Support:** The trial was funded by the British Heart Foundation (SP/12/10/29922 and PG/15/51/31596). Dr Mills is supported by the Butler Senior Clinical Research Fellowship (FS/16/14/32023), and Drs Lee and Wereski are supported by a Clinical Research Training Fellowship (FS/18/25/33454) and a Research Excellence Award (RE/18/5/34216), respectively, from the British Heart Foundation. Dr Chapman receives support from a Starter Grant for Clinical Lecturers by the Academy of Medical Sciences (SGL021/1075).

**Role of the Funder/Sponsor:** The funding sources had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

**Additional Contributions:** We thank the High-STEACS investigators for their contributions. **Chief investigator:** Dr Mills; *University of Edinburgh*. **Trial managers:** Dr Fiona E Strachan; *University of Edinburgh* and Mr Christopher Tuck; *Edinburgh Clinical Trials Unit*. **Trial research team:** Dr Anoop S V Shah, Dr Fiona E Strachan, Dr Atul Anand, Dr Anda Bularga, Dr Wereski, Dr Amy V Ferry, Dr Lee, Dr Chapman, Mr Dennis Sandeman, Dr Philip D Adamson, Dr Catherine L Stables, Dr Catalina A Vallejo, Dr Athanasios Tsanasis, Ms Lucy Marshall, Ms Stacey D Stewart, Dr Takeshi Fujisawa, Ms Mischa Hautvast, Ms Jean McPherson, and Ms Lynn McKinlay; *University of Edinburgh*. **Grant applicants:** Dr Mills, Prof David E Newby, Prof Keith AA Fox, Dr Simon Walker, and Dr Christopher J Weir; *University of Edinburgh*; Prof Colin Berry; *University of Glasgow*. **Adjudication panel:** Dr Anoop S V Shah, Dr Atul Anand, Dr Chapman, Dr Lee, Dr Jack Andrews, Dr Phil Adamson, Dr Alastair Moss, Dr Mohamed Anwar, Dr John Hung, and Dr Mills; *University of Edinburgh*. **Trial steering committee:** Dr Mills, Prof David Newby, Prof Dr Gray, Prof Keith AA Fox, Dr Simon Walker, Prof John Norrie, Prof Christopher Weir; *University of*

Edinburgh; Prof Colin Berry, Prof Ian Ford, Dr David A McAllister; *University of Glasgow*; Prof Paul O Collinson; *St George's University Hospitals*; Prof Fred S Apple; *University of Minnesota*; Mr Alan Reid; *UKNEQAS*; Dr Anne Cruikshank, Dr Iain Findlay, Dr Donogh Maguire; *NHS Greater Glasgow and Clyde*; Dr Shannon Amoils; *British Heart Foundation*, Ms Jennifer Stevens. **Biochemistry sub-group committee:** Dr Simon Walker; *University of Edinburgh*; Dr Jonathan Malo, *NHS Lothian*; Mr Alan Reid; *UKNEQAS*; Dr Anne Cruikshank; *NHS Greater Glasgow and Clyde*; and Prof Paul O Collinson; *St George's University Hospitals*; **Data monitoring committee:** Prof Colin Fischbacher; *Public Health Scotland*; Dr Bernard Croal; *NHS Grampian*; and Prof Stephen J Leslie; *NHS Highland*. **Edinburgh Clinical Trials Unit:** Ms Catriona Keerie, Prof Christopher Weir, Mr Richard Parker, Mr Allan Walker, Mr Ronnie Harkess, Mr Chris Tuck, and Mr Tony Wackett. **NHS Greater Glasgow & Clyde Safe Haven:** Dr Roma Armstrong, Ms Marion Flood, Ms Laura Stirling, Ms Claire MacDonald, Mr Imran Sadat, and Mr Frank Finlay. **NHS Lothian eHealth and Safe Haven:** Dr Heather Charles, Ms Pamela Linksted, Mr Stephen Young, Mr Bill Alexander, and Mr Chris Duncan. The High-STEACS investigators were responsible for the conception and design of the High-STEACS trial, and the acquisition of data used in this analysis.

1. Chapman AR, Lee KK, McAllister DA, et al. Association of high-sensitivity cardiac troponin I concentration with cardiac outcomes in patients with suspected acute coronary syndrome. *JAMA*. 2017;318(19):1913-1924. doi:10.1001/jama.2017.17488
2. Bularga A, Lee KK, Stewart S, et al. High-sensitivity troponin and the application of risk stratification thresholds in patients with suspected acute coronary syndrome. *Circulation*. 2019;140(19):1557-1568. doi:10.1161/CIRCULATIONAHA.119.042866
3. Thygesen K, Alpert JS, Jaffe AS, et al. Fourth universal definition of myocardial infarction (2018). *Eur Heart J*. 2019;40(3):226. doi:10.1093/eurheartj/ehy856
4. Roffi M, Patrono C, Collet JP, et al; ESC Scientific Document Group. 2015 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation: task force for the Management of Acute Coronary Syndromes in Patients Presenting without Persistent ST-Segment Elevation of the European Society of Cardiology (ESC). *Eur Heart J*. 2016;37(3):267-315. doi:10.1093/eurheartj/ehv320
5. Shah ASV, Anand A, Strachan FE, et al; High-STEACS Investigators. High-sensitivity troponin in the evaluation of patients with suspected acute coronary syndrome: a stepped-wedge, cluster-randomised controlled trial. *Lancet*. 2018;392(10151):919-928. doi:10.1016/S0140-6736(18)31923-8
6. Chapman AR, Adamson PD, Shah ASV, et al; High-STEACS Investigators. High-sensitivity cardiac troponin and the universal definition of myocardial infarction. *Circulation*. 2020;141(3):161-171. doi:10.1161/CIRCULATIONAHA.119.042960

## COMMENT & RESPONSE

### Interpreting the Need for Implantable Loop Recorder Monitoring in Pregnant Women at High Risk of Arrhythmias

**To the Editor** We congratulate Sliwa et al<sup>1</sup> for their excellent analysis on the use of implantable loop recorders (ILRs) in addition to 24-hour Holter electrocardiography to improve the detection of arrhythmias in pregnant women with cardiovascular disease compared with 24-hour Holter electrocardiography alone. We believe these findings to be pertinent, as a 2019 study by Moulig et al<sup>2</sup> observed 66 patients with peripartum cardiomyopathy over 5 years and found that 17% had arrhythmias, including paroxysmal supraventricular tachycardia, ventricular tachycardia, or ventricular fibrillation. It should be noted the large number of arrhythmias found in this population may be because 9 patients had wearable cardioverter-defibrillators inserted at diagnosis. Of these patients, 8 also had permanent implantable cardioverter-defibrillators inserted.<sup>2</sup> These data show that patients with peripartum cardiomyo-

pathy remain at high risk of arrhythmias for prolonged periods and could see considerable benefit from monitoring as described by Sliwa et al.<sup>1</sup>

Furthermore, Cuneo et al<sup>3</sup> looked at 148 pregnancies from families with long QT syndromes. They found that stillbirths (fetal death after 20 weeks' gestation) were far more frequent in those with long QT syndrome compared with the general population (4% vs approximately 0.5%),<sup>3</sup> making close monitoring with ILRs in high-risk populations even more pertinent. Another study by Schlichting et al<sup>4</sup> studied 17 729 pregnancies in women with congenital heart disease. They found that the risk of arrhythmias was significantly increased (adjusted odds ratio, 12.4; 95% CI, 11.0-14.0) and that ILRs should be considered in this population.

Li et al<sup>5</sup> have explored treatment of maternal arrhythmias and showed in a series of 28 pregnant patients that catheter ablation for sustained maternal tachyarrhythmias using intracardiac echocardiography catheter and zero fluoroscopy is a viable technique. This makes the data from Sliwa et al<sup>1</sup> more relevant as there is an option for treatment for some of these women. On the background of these results, it stands that ILR monitoring in high-risk populations may soon become a necessity to improve fetal outcomes and further points to the need for research into the management of pregnant women with arrhythmias.

**Syed Yousaf Ahmad**  
**Awais Ahmad, MD**  
**Falaq Syed Raheel, MPhil**

**Author Affiliations:** Faculty of Medicine, Imperial College London, London, United Kingdom (S. Y. Ahmad); Faculty of Medicine and Surgery, University of Malta, Msida, Malta (A. Ahmad); College of Medical and Dental Services, University of Birmingham, Birmingham, United Kingdom (Raheel).

**Corresponding Author:** Syed Yousaf Ahmad, Faculty of Medicine, Imperial College London, Level 2, Faculty Building, South Kensington Campus, London SW7 2AZ, United Kingdom (syed.ahmad17@imperial.ac.uk).

**Published Online:** August 26, 2020. doi:10.1001/jamacardio.2020.3521

**Conflict of Interest Disclosures:** None reported.

1. Sliwa K, Azibani F, Johnson MR, et al. Effectiveness of implanted cardiac rhythm recorders with electrocardiographic monitoring for detecting arrhythmias in pregnant women with symptomatic arrhythmia and/or structural heart disease: a randomized clinical trial. *JAMA Cardiol*. 2020;5(4):458-463. doi:10.1001/jamacardio.2019.5963
2. Moulig V, Pfeffer TJ, Ricke-Hoch M, et al. Long-term follow-up in peripartum cardiomyopathy patients with contemporary treatment: low mortality, high cardiac recovery, but significant cardiovascular co-morbidities. *Eur J Heart Fail*. 2019;21(12):1534-1542. doi:10.1002/ehf.1624
3. Cuneo BF, Kaizer AM, Clur SA, et al; Fetal LQTS Consortium. Mothers with long QT syndrome are at increased risk for fetal death: findings from a multicenter international study. *Am J Obstet Gynecol*. 2020;222(3):263.e1-263.e11.
4. Schlichting LE, Insaft TZ, Zaidi AN, Lui GK, Van Zutphen AR. Maternal comorbidities and complications of delivery in pregnant women with congenital heart disease. *J Am Coll Cardiol*. 2019;73(17):2181-2191. doi:10.1016/j.jacc.2019.01.069
5. Li MM, Sang CH, Jiang CX, et al. Maternal arrhythmia in structurally normal heart: prevalence and feasibility of catheter ablation without fluoroscopy. *Pacing Clin Electrophysiol*. 2019;42(12):1566-1572. doi:10.1111/pace.13819

**In Reply** We thank Ahmad and colleagues for their letter regarding our article.<sup>1</sup> Evidence for best practices in the diagnosis and management of arrhythmias in pregnant and peripartum